



ICD-10-PCS Code Request Meniscus Replacement with the NUsurface[®] Synthetic Meniscus Implant

ICD-10-PCS C&M Meeting – Sept. 14, 2021



Clinical Presenter: Wayne Gersoff, MD, Orthopedic Surgery and Sports Medicine
Advanced Orthopedics & Sports Medicine Specialists, Denver, CO

- NUsurface SUN Clinical Trial Investigator
- Medical Education Consultant – *Active Implants, LLC*

Clinical Presenter: Tom Carter, MD, Orthopedic Surgery and Sports Medicine
Banner University, Professor of Orthopedic Surgery, Phoenix, AZ

- NUsurface SUN Clinical Trial Investigator
- Medical Education Consultant – *Active Implants, LLC*

Company Representative: Erik Harris, MHA, VP Global Market Access & Reimbursement
Active Implants, LLC




Overview of the Meniscus, Incidence of Injury and Outcomes


- The **meniscus protects the knee by distributing loads** from the femoral condyles to the tibial plateau, absorbing shock during motion, and reducing friction between the articulating bones¹
- **Meniscus tears** can result at any time from excessive forces caused by **knee trauma** (*acute tears are more common in the 20s-30s age bands*), or the result of natural, **age-related degeneration, resulting in meniscus dysfunction** (*degenerative tears are more common in the 40s-60s age bands*)²
 - ❖ *Medial meniscus tears are up to 5x more frequent than lateral meniscus tears*³
- **Over 760,000 partial meniscectomies are performed each year**, making it the most common surgical treatment for meniscus-related knee pain and/or loss of knee function⁴
- **25% of patients (190,000) treated with partial meniscectomy have persistent symptoms within 2 years after surgery**, and 40% (304,000) within 10-15 years⁵⁻⁸
 - ❖ *Partial meniscectomy patients have a 10x higher likelihood for a knee replacement*⁹
- **Non-surgical treatments** of meniscus dysfunction **are ineffective** in the long-term, do not maintain efficacy over repeated procedures, and **do not address the underlying disease state**¹⁰⁻¹²

A Damaged / Dysfunctional Meniscus Leads to Increased Femur to Tibia Load Concentration and Eventual Cartilage Loss and Pain


Intact Meniscus




Intact Meniscus
Pressure
Distribution¹³




Torn Meniscus




Post Meniscectomy



Concentrated
Focal Pressure
Distribution¹³




← Femur to Tibia contact
pressure map →



- Partial Meniscectomy Increases Risk of Knee Replacement by **more than 10 times**⁹
- Younger Patient Age for Meniscectomy Increases Risk of Knee Replacement **by Nearly 40 Times**⁵⁻⁸
- In a British longitudinal study, by 15 years After Meniscectomy **13.5% of Patients Had Knee Replacements**¹⁴


Meniscus Dysfunction – Current Treatment Options

There is a substantial unmet need for a product that can reduce pain and enable function, thereby bridging the many years between meniscectomy/repair and eventual knee replacement




The NUsurface® Meniscus Implant was clinically studied to treat patient pain and function in the “gap” between current meniscus surgical treatment* and an irreversible knee joint replacement


How Does the NUsurface® Implant Work



Mechanism of Action



NUsurface® Restores Knee Pressure Distribution Similar to an Intact, Articulating Meniscus^{13, 15}



***NUsurface® manufactured in 7 sizes in Left and Right configurations**

(Note: final number of sizes TBD per FDA market authorization)


Surgical Technique – SIMPLIFIED Procedure Overview

Vignette and Procedure Fully Described in the Submitted ICD-10-PCS Code Request

- Pre-operative sizing via MRI assessment (*uses existing patient diagnostic MRI*)
- **Procedure may be completed in approximately 1-1.5 hours** (*per surgeon experience and case complexity*)
- 6 Week Rehab program, weightbearing same day


1

Prep & Drape for Procedure




2

Arthroscopic Assisted Joint Preparation



3

**Open Arthrotomy
NUsurface® Implantation**




4


**NUsurface®
Final Position**

**Medial
Meniscus Rim**

**NUsurface®
Meniscus
Implant**



- **242 Subject MERCURY Clinical Dataset** is currently Under FDA De Novo 510(k) Evaluation comprising:
 - ✓ **VENUS¹⁶ = 127 Subject** Randomized Clinical Trial NUsurface (N=61) to Non-Surgical Controls (N=66)
 - ✓ **SUN¹⁷ = 115 Subject** Prospective Longitudinal Single-Arm Interventional Trial
- Endpoints for **KOOS Pain** and **KOOS Overall** of **20-point improvement at 24 months**
 - ❖ Inclusion/Exclusion criteria identified in the ICD-10-PCS code request
- **FDA Breakthrough Device Designation – Sept. 13, 2019**




MERCURY = VENUS RCT + SUN Single-Arm Across 23 Clinical Trial Sites

Richard Alfred, MD (Capital Region Orthopedics – Albany, NY)
Maxwell Alley, MD (Capital Region Orthopedics – Albany, NY)
Jack Farr, MD (OrthoIndy – Indianapolis, IN)
William Garrett, MD (Duke University MC – Raleigh, NC)
Thomas Giel, MD (OrthoMemphis – Memphis, TN)
Andreas Gomoll, MD (Hospital for Special Surgery, New York, NY)
Elliott Hershman, MD (North Shore LIJ, Lenox Hill Hospital – New York, NY)
Randall Holcomb, MD (OrthoMemphis – Memphis, TN)
Christopher Kaeding, MD (Wexner Medical Center, OSU – Columbus, OH)
Christian Lattermann, MD (Brigham and Women's Hospital – Boston, MA)
Brian McKeon, MD (New England Baptist Hospital – Boston, MA)
Claude Moorman, MD (Duke University MC – Raleigh, NC)
Allison Toth, MD (Duke University MC – Raleigh, NC)
Kenneth Zaslav, MD (OrthoVirginia – Richmond, VA)


Larry Bankston, MD (BRORTH – Baton Rouge, LA)
Joseph Berman, MD (Baylor Orthopedic and Spine Hospital - Dallas, TX)
Thomas Carter, MD (TOCA – Phoenix, AZ)
Andrew Cooper, MD (Comprehensive Orthopedics – Salt Lake City, UT)
Robert Easton, MD (BRORTH – Baton Rouge, LA)
Richard Edelson, MD (Oregon Sports Medicine – Portland, OR)
Rachel Frank, MD (CU Sports Medicine – Denver, CO)
Wayne Gersoff, MD (AdvancedOrtho – Denver, CO)
Jonathan Greenleaf, MD (Oregon Sports Medicine – Portland, OR)
Scott Hacker, MD (Horizon Clinical Research, San Diego, CA)
Deryk Jones, MD (Ochsner Sports Medicine – New Orleans, LA)
Peter Kurzweil, MD (Memorial Orthopaedics – Long Beach, CA)
Eric McCarty, MD (CU Sports Medicine -Boulder, CO)
William Montgomery, MD (St. Mary's Medical Center - San Francisco, CA)
Noah Weiss, MD (Weiss Orthopedics, Sonoma, CA)

VENUS TRIAL

- Randomized
- 61 NUsurface® Implant Patients
- 66 Control Patients
- 10 sites



- FDA-Pooled Randomized Data
- 176 NUsurface® Implant Patients
- 66 Control Patients




SUN TRIAL

- Single Arm
- 115 NUsurface® Implant Patients
- 13 sites

Clinical Outcomes • NUsurface® KOOS Pain by Stratified Age Bands

Responder Rates at Different Degrees of KOOS Pain Improvement

Baseline Age Range	Total	No Improvement (%)	> 10 points (%)	> 20 points (%)	>30 points (%)
30's	27	14.8%	85.1%	63%	40.7%
40's	45	20%	80%	71.1%	46.6%
50's	51	17.6%	82.4%	64.7%	43.1%
60's	30	6.7%	93.3%	76.6%	46.6%



NOTE: 12-month trial observations have been published^{18, 19} while the 24-month data is currently under FDA review and has not yet been published

Clinical Trial Patient Age Demographics

It's not the Age of the Patient but the "Age" of the Patient's Knee


Trial	N Total	Average Age	Standard Deviation	Oldest Treated Patient	Youngest Treated Patient	Total Patients >63 yrs	% >63 yrs at Enrollment	Total Patients ≥65 yrs	% ≥65 yrs at Enrollment
MERCURY	242	49.8	10.1	69	30	23	9.5%	18	7.4%
VENUS	127	50.4	10.7	69	30	14	11%	10	7.9%
SUN	115	49.2	9.4	69	30	9	7.9%	8	7.0%

- Clinical Trial Population Represents a Generalizable Population:
 - Of the **9.5%** treated subjects who were ≥ 65 years old During the 2-year trial
 - **7.4%** were ≥ 65 at the Start of the clinical trial

NOTE: 12-month trial observations have been published^{18, 19} while the 24-month data is currently under FDA review and has not yet been published

Clinical Outcomes KOOS Pain • Statistically Superior Efficacy Results

KOOS_{PAIN}: Non-Surgical Control vs MERCURY Investigational



Data Locked 30 June 2020. Error bars represent the 95% CI


Sample size: **MERCURY** (Baseline n=176, 1.5 months n=175, 6 months n=167, 12 months n=165, 24 months n=153)

Sample size: **Control** (Baseline n=66, 1.5 months n=63, 6 months n=56, 12 months n=46, 24 months n=43)

NOTE: 12-month trial observations have been published^{18, 19} while the 24-month data is currently under FDA review and represents FDA-submitted data on file (i.e., pre-publication)

Clinical Outcomes KOOS Overall • Statistically Superior Efficacy Results

KOOS OVERALL: Non-Surgical Control vs MERCURY Investigational



Data Locked 30 June 2020. Error bars represent the 95% CI
Sample size: **MERCURY** (Baseline n=176, 1.5 months n=166, 6 months n=167, 12 months n=165, 24 months n=153)
Sample size: **Control** (Baseline n=66, 1.5 months n=62, 6 months n=56, 12 months n=46, 24 months n=43)

NOTE: 12-month trial observations have been published^{18, 19} while the 24-month data is currently under FDA review and represents FDA-submitted data on file (i.e., pre-publication)


The NUsurface® Implant Maintained Cartilage while Reducing Pain

Measurement of Knee Cartilage Condition in Control Subjects vs. NUsurface® Subjects at Baseline and at 2-Years


% Full Thickness Cartilage Lesion on Distal Medial Femoral Condyle

Full Thickness Cartilage Lesion Exposing Bone ■ YES ■ NO

Cartilage **Degeneration** in Control Subjects



Cartilage **Stability** in NUsurface® Subjects



NOTE: 12-month trial observations have been published^{18, 19} while the 24-month data is currently under FDA review and has not yet been published

NUsurface® Meniscus Implant

2021 Pre-FDA • Corporate Communication

CMS ICD-10-PCS C&M Committee Presentation


Not for distribution in the United States and its territories.

ACTIVE IMPLANTS®
Copyright ©2021 Active Implants, LLC. All Rights Reserved.

CAUTION – Investigational Device. Limited by U.S. law to investigational use.

Patient Clinical Trial Accountability in the NUsurface® Group

90% of NUsurface® Subjects Completed 24 Months Compared with 65% of Control Subjects



NOTE: 12-month trial observations have been published^{18, 19} while the 24-month data is currently under FDA review and has not yet been published

NUsurface® Meniscus Implant

2021 Pre-FDA • Corporate Communication
CMS ICD-10-PCS C&M Committee Presentation

Not for distribution in the United States and its territories.

ACTIVE IMPLANTS®
Copyright ©2021 Active Implants, LLC. All Rights Reserved.

CAUTION – Investigational Device. Limited by U.S. law to investigational use.

Site of Service

The NUsurface® meniscus implant surgeries in the clinical trials were performed in the outpatient hospital and ASC settings. Because the anticipated NUsurface® patient age range in an anticipated FDA label is likely to be similar to that of the clinical trials (i.e., ages 30-75), it is anticipated that in general use post FDA-decision that surgery in an inpatient setting may be performed dependent on patient comorbidities

Hospital Outpatient Surgery

NUsurface® meniscus implant surgery performed in a hospital outpatient setting or an ASC setting

Outpatient Ambulatory Surgery Center



Inpatient Setting

NUsurface® meniscus implant surgery may be performed during an inpatient stay (low incidence), dependent on patient comorbidities

References

1. Fox, A.J., A. Bedi, and S.A. Rodeo, The basic science of human knee menisci: structure, composition, and function. *Sports Health*, 2012. 4(4): p. 340-51.
2. Fox, A.J., et al., The human meniscus: a review of anatomy, function, injury, and advances in treatment. *Clin Anat*, 2015. 28(2): p. 269-87.
3. Lento, P.H. and V. Akuthota, Meniscal injuries: A critical review. *J Back Musculoskelet Rehabil*, 2000. 15(2): p. 55-62.
4. Hall, M.J., et al., Ambulatory Surgery Data From Hospitals and Ambulatory Surgery Centers: United States, 2010, U.S.D.o.H.a.H. Services and C.f.D.C.a. Prevention, Editors. 2017: Natl Health Stat Report. p. 1-15.
5. Sihvonen, R., et al., Arthroscopic partial meniscectomy versus placebo surgery for a degenerative meniscus tear: a 2-year follow-up of the randomised controlled trial. *Ann Rheum Dis*, 2018. 77(2): p. 188-195.
6. Yim, J.H., et al., A comparative study of meniscectomy and nonoperative treatment for degenerative horizontal tears of the medial meniscus. *Am J Sports Med*, 2013. 41(7): p. 1565-70.
7. Katz, J.N., et al., Surgery versus physical therapy for a meniscal tear and osteoarthritis. *N Engl J Med*, 2013. 368(18): p. 1675-84.
8. Chatain, F., et al., The natural history of the knee following arthroscopic medial meniscectomy. *Knee Surg Sports Traumatol Arthrosc*, 2001. 9(1): p. 15-8.
9. Abram, S.G.F., et al., Long-term rates of knee arthroplasty in a cohort of 834 393 patients with a history of arthroscopic partial meniscectomy. *Bone Joint J*, 2019. 101-B(9): p. 1071-1080.
10. Filardo, G., et al., No Effects of Early Viscosupplementation After Arthroscopic Partial Meniscectomy: A Randomized Controlled Trial. *Am J Sports Med*, 2016. 44(12): p. 3119-3125.

References (continued)

11. McAlindon, T.E., et al., Effect of Intra-articular Triamcinolone vs Saline on Knee Cartilage Volume and Pain in Patients With Knee Osteoarthritis: A Randomized Clinical Trial. *JAMA*, 2017. 317(19): p. 1967-1975.
12. Kompel, A.J., et al., Intra-articular Corticosteroid Injections in the Hip and Knee: Perhaps Not as Safe as We Thought? *Radiology*, 2019. 293(3): p. 656-663.
13. Linder-Ganz E, et al., A novel quantitative approach for evaluating contact mechanics of meniscal replacements. *J Biomech Eng*. 2010 Feb;132(2):024501
14. Abram, S.G.F, et al., Long-term rates of knee arthroplasty in a cohort of 834 393 patients with a history of arthroscopic partial meniscectomy. *Bone Joint J*, 2019; 101-B:1071-1080
15. Shemesh M, Shefy-Peleg A, Levy A, Shabshin N, Condello V, Arbel R, Gefen A. Effects of a Novel Medial Meniscus Implant on the Knee Compartments: Imaging and Biomechanical Aspects. *Biomechanics and Modeling in Mechanobiology*, 2020 March 31
16. ClinicalTrials.gov: NCT02108496
17. ClinicalTrials.gov: NCT02483988
18. McKeon BP et al., Preliminary Results From a US Clinical Trial of a Novel Synthetic Polymer Meniscal Implant. *Orthop J Sports Med*. 2020 Sep 29;8(9).
19. Zaslav K, et al., Treatment of post-meniscectomy knee symptoms with medial meniscus replacement results in greater pain reduction and functional improvement than non-surgical care. *Knee Surg Sports Traumatol Arthrosc*. 2021 Apr 21

Bionate®
Thermoplastic
Polycarbonate-
urethane (PCU)



UHMWPE* Fibers



Anterior (A)

Lateral (L)

Medial (M)



Posterior (P)




Anterior



Posterior

*Ultra-high molecular weight polyethylene

*(Note: final Product & Instruments TBD per FDA market authorization)



1. Meniscus Probe

2. Universal Insertion
Instrument

3. Extraction Instrument

4. Bone Rasp, Right + Left

